

CCM.0000000000006253. Epub 2024 Mar 20. PMID: 38506571.

13. Giani M, Rezoagli E, Barbaro RP, Riera J, Bellani G, Brochard L, et al. Noninvasive Ventilation Before Intubation and Mortality in Patients Receiving Extracorporeal Membrane Oxygenation for COVID-19: An Analysis of the Extracorporeal Life Support Organization Registry. *ASAIO J.* 2024 Jul 1;70(7):633-639. doi: 10.1097/MAT.0000000000002132. Epub 2024 Jun 27.

14. Fossali T, Locatelli M, Colombo R, Veronese A, Borghi B, Ballone E, et al. Awake pronation with helmet CPAP in early COVID-19 ARDS patients: effects on respiratory effort and distribution of ventilation assessed by EIT. *Intern Emerg Med.* 2024 Oct;19(7):2025-2034. doi: 10.1007/s11739-024-03572-0. Epub 2024 Mar 26. PMID: 38532048.

15. Liu L, Sun Q, Zhao H, Liu W, Pu X, Han J, et al; Chi-ARDS Net (Chinese ARDS Research Network). Prolonged vs shorter awake prone positioning for COVID-19 patients with acute respiratory failure: a multicenter, randomised controlled trial. *Intensive Care Med.* 2024 Aug;50(8):1298-1309. doi: 10.1007/s00134-024-07545-x. Epub 2024 Aug 1. PMID: 39088076; PMCID: PMC11306533.

16. González-Castro A, Huertas Martín C, Cuenca Fito E, Peñasco Y, Gonzalez C, Rodríguez Borregán JC. Duration of the first prone positioning maneuver and its association with 90-day mortality in patients with acute respiratory failure due to COVID-19: A retrospective study of time terciles. *Med Intensiva (Engl Ed).* 2024 Aug;48(8):457-466. doi: 10.1016/j.medine.2024.04.009. Epub 2024 Apr 30. PMID: 38688818.